

**Exercises**

E22.17 [1RK] Discuss the differential equation


$$\begin{cases} y'(x) = \frac{1}{y(x)-x^2} \\ y(0) = a \end{cases}$$

for  $a \neq 0$ , studying in a qualitative way the existence (local or global) of solutions, and the properties of monotonicity and convexity/concavity.<sup>a</sup>

Show that the solution exists for all positive times.


Show that for  $a > 0$  the solution does not extend to all negative times.

*Difficulty:\**. Show that there is a critical  $\tilde{a} < 0$  such that, for  $\tilde{a} < a < 0$  the solution does not extend to all negative times, while for  $a \leq \tilde{a}$  the solution exists for all negative times; also for  $a = \tilde{a}$  you have  $\lim_{x \rightarrow -\infty} y(x) - x^2 = 0$ .



In dotted purple the line of inflections. In dashed red the parabola where the derivative of the solution is infinite. In yellow the solutions with initial data  $y(0) = 2, y(0) = 1, y(0) = 1/1000$ .

Figure 9: Exercise 22.17. Solutions for  $a > 0$



In dotted purple the line of inflections. In dashed red the parabola where the derivative of the solution is infinite. Solutions are drawn with initial data  $a = -1.4$  ("green"),  $a = -1.0188$  ("orange") and  $a = -1.019$  ("yellow"). Note that the latter two differ only by 0.0002 in their initial data (indeed they are indistinguishable in the graph for  $x > -1$ ), but then for  $x < -1$  they move apart quickly, and for  $x = -2$  they are respectively 3.25696 and 2.54856, with a difference of about 0.7!

Figure 10: Exercise 22.17. Solutions for  $a < 0$

**Solution 1.** [1RP]

<sup>a</sup>The differential equation is taken from exercise 13 in [2].